

Epigenetic mechanisms and associated brain circuits in the regulation of positive emotions: a role for Transposable Elements.

Simona Gaudi ¹, Guia Guffanti ², James Fallon³, Fabio Macciardi ^{3, 4, 5, 6}

¹ Department of Infectious, Parasitic and Immune-Mediated Diseases, Italian National Institute of Health, Rome, Italy;

² Department of Psychiatry, Columbia University, New York, New York

³ Department of Psychiatry and Human Behavior, University of California Irvine, Irvine, California, USA

⁴ Center for Autism Research and Treatment (CART), University of California, Irvine, California, USA

⁵ Center for Epigenetics and Metabolism, University of California, Irvine, California, USA

⁶ Department of Pharmacological and Bio-molecular Sciences, University of Milano, Italy

Abbreviated title: Transposable Elements and the emotional brain

Key-words: brain circuits, positive emotions, genomics, epigenetics Transposable Elements

Corresponding Author:

Fabio Macciardi

Department of Psychiatry and Human Behavior

School of Medicine

University of California, Irvine

5251 California Ave, Suite 240

Irvine, CA 92617

Telephone: 949-824-4559

Fax: 949-824-3324

Email: fmacciar@uci.edu

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as an 'Accepted Article', doi: 10.1002/cne.24046

© 2016 Wiley Periodicals, Inc.

Received: Mar 18, 2015; Revised: May 23, 2016; Accepted: May 23, 2016

This article is protected by copyright. All rights reserved.

Abstract (150 words)

Epigenetic programming and reprogramming are at the heart of cellular differentiation and represent developmental and evolutionary mechanisms both in germ-line and somatic cell lines. Only about 2 % of our genome is composed of protein-coding genes, while the remaining 98 %, once considered “junk” DNA, codes for regulatory / epigenetic) elements that control how genes are expressed in different tissues and across time from conception to death. While we already know that epigenetic mechanisms are at play in cancer development and in regulating metabolism (cellular and whole body), the role of epigenetics in the developing prenatal and postnatal brain, and in maintaining a proper brain activity throughout the various stages of life, in addition to having played a critical role in human evolution, is a relatively new domain of knowledge. Here we present the current state-of-the-art techniques and results of these studies within the domain of emotions, and then speculate how genomic and epigenetic mechanisms can modify and potentially alter our emotional, limbic, brain and affect our social interactions.

Introduction

The completion of the human genome sequencing has represented the beginning of a genetic and genomic revolution in health care. The pace of genomic medicine has moved at a surprisingly fast rate. Within 70 years we have progressed from the foundational studies of Friedrich Miescher, Phoebus Levene, and Erwin Chargaff, and then Watson and Crick's seminal discovery of the double helix, to reading the whole genetic material in a human being with the possibility of using the genomic information for medical and health-related applications. However, and despite having obtained the complete sequence of our genome, we still need substantial new research to really understand the functional significance of the genome architecture, particularly within the context of neuroscience. While we have been developing a better knowledge of the most fundamental molecular mechanisms that regulate neuronal functions, recently we have in parallel, been acquiring a better understanding of brain circuits and functions; the issue at stake is how to integrate these disparate information into a coherent framework.

Our understanding of the structural and functional composition of the genome, and the plasticity of gene regulatory pathways has increased exponentially, both through technical developments and the resultant progressive understanding of common and complex diseases. Unexpectedly, however, the completion of the mapping and characterization of the human genome has revealed that the true complexity of the genome has little to do with the simple number, and even variability, of its coding genes. Only 2% of the human genome codes for proteins; the remaining 98% is represented by non-protein coding sequences. For decades the non-coding genome was considered "junk" DNA without any functional significance, but deep sequencing analyses in recent years suggest that the whole human genome is **NOT** transcribed into RNAs (Dunham et al., 2012; [Harrow et al., 2012](#); Kundaje et al., 2015). The emerging pattern is that the vast

majority of our genome is functional and regulates the gene-coding machinery by producing non-coding RNAs in a myriad of ways. Another level of complexity is related to the epigenetic interplay between genome activity and environmental influences: current research into functional genomics and individual deep sequencing are opening new perspectives in understanding how genome and environment interact in complex diseases ([Coop et al., 2010](#)).

Here we present a short review of our current knowledge of the epigenetic mechanisms that functionally regulate the genome most involved in regulation of the central nervous system and which contribute to the proper, context-dependent functions of brain circuits involved in positive (or negative) emotions also via external (i.e., non-genomic) stimuli. For example, it is well known that social adversities, physical abuse, loneliness, and even grief are associated with increased susceptibility to disease. When such stressful events occur, they stimulate an adapted behavioral response where the environmental challenges can be viewed within the framework of the evolutionary-based adaptive match/mismatch hypothesis (Schmidt, 2011; [Daskalakis et al., 2012](#); [Daskalakis et al., 2013](#)). Under this scenario, the outcome of a stressful condition leading to either a physiological resilience or to a psychopathological condition, depends on the coping strategies that any individual has developed when faced with early life events. Coping strategies emerge as the result of the interaction between the “programmed” genomic-based adaptations with the constraints due to early, adverse events, especially in the perinatal and postnatal periods.

Molecular architecture of the emotional brain: what do we really know?

A vast body of literature in the last decade attests to the renewed and increased interest in the study of the neural bases of affections or emotions. New analytical strategies that combine refined brain imaging and genomic methods, together with psychometric and clinical

quantitative traits, are allowing for the current research in neuroscience to clarify how our brain and our behavior are shaped both phylogenetically and evolutionarily. As thoroughly reviewed by Panskepp and Biven (2012) the “emotional” (primarily composed of the limbic system) and the “social” brain (primarily composed of the prefrontal and anteromedial temporal systems) are privileged targets for these studies, but they represent more difficult objectives than, say, the more accessible and long-studied cognitive and executive functions of the dorsal and lateral neocortical systems. To explore the world of human emotions, we need both theoretical and operational improvements. Theoretically, we have already made substantial improvements, shifting from a purely psychological to a more integrated neurobiological / cognitive framework that can help evince a clearer understanding of experimental findings. Operationally, sophisticated investigations are required to understand how genomic and epigenomic mechanisms that regulate brain circuits controlling for emotions and social behaviors work day to day and over the full lifetime in living humans. While other papers in this special issue present in details the current hypotheses of brain circuits and neurotransmitter pathways that control for positive emotions, here we are interested to speculate on a model that integrates the functional and the molecular architecture of the emotional brain.

In short, we know that both subcortical and cortical structures are relevant in controlling for affects and emotions in humans, including the orbitofrontal cortex, the amygdala, the hippocampus, the limbic system and their associated cortico-subcortical loops in the basal ganglia, limbic thalamus and brainstem monoaminergic nuclei. Our knowledge of the functional anatomy of these brain regions has steadily progressed in the last 30 years, but other brain regions also considered relevant, like the periaqueductal grey, the septum and a few non-monoaminergic brainstem nuclei are still poorly known. Moreover, little is known about the developmental mechanisms that control for the proper circuital organization of these structures

and of their neuronal connections. We know since quite a long time the essential role played by dopamine in the “reward” system ([Schultz, 2011](#); [Russo and Nestler, 2013](#)) and the key role played by other neurotransmitters like the opioid peptides and endocannabinoids. Experimental support for the relevance of these systems is increasing: for example, a frequent DNA variant in the FAAH gene that reduces the amount of deactivation of anandamide, a naturally occurring human cannabinoid, correlates with lower levels of anxiety and fear. The reduced FAAH expression associated with the variant allele also “selectively enhances prefronto-amygdala connectivity and fear extinction learning [...] suggest[ing] a gain of function in fear regulation” ([Dincheva et al., 2015](#)). This mechanism may represent an adaptive function for a primary process of positive emotion that humans share with other mammals, in this case mice.

It is possible, however, that different mutational mechanisms other than nucleotide substitutions must be invoked to explain how gene networks have been (re)wired during human evolution to deal with environmental, primarily intraspecies control of emotions and hence have shaped our emotional and social brain. Widespread epigenetic changes have been proposed to underlie alterations in gene expression associated with gene networks and brain circuits that control for positive emotions. Modular gene-networks that are co-expressed in anatomically different, but functionally related brain regions, can represent the backbone of a circuital organization that modulates our ability to deal with emotions. Under this scenario, genes that belong to the same network or genes that follow a hierarchical time-dependent expression developmental pattern must be activated in a coordinated way across different and somewhat distant brain regions. This coordinated molecular / functional activity across brain regions is similar to the mechanisms that are putatively at work in complex neuropsychiatric disorders, including schizophrenia and affective disorders, i.e. in all cases when scores of genes must be jointly implicated epistatically.

However, how both normal and pathological neuronal circuitry depends on the specificity and extent of epigenetic programs has not yet been fully elucidated.

New insights from genomic studies.

One of the surprises of the genomic era is the relative small number of protein-coding genes in human genome. Before the completion of the Human Genome Project indirect estimates for the number of protein-coding genes were ranging from 35,000 to 100,000 ([Ohno, 1972](#); [Lander et al., 2001](#)). To date, the number of protein-coding genes has been experimentally reduced to 20-25,000, and we expect it to be further revised (Consortium, 2004). Current evidence shows that many regulatory-complexes are needed in the development and function of the vertebrate brain ([Yoo and Crabtree, 2009](#); [Yoo et al., 2009](#)): recent progress from the ENCODE project has begun to reveal the complexities of the non-coding human genome, beginning with the functional characterization of an unprecedented number of elements previously considered non-functional ([Kundaje et al., 2015](#)). These efforts have lead to the generation of an extensive catalog of more than four million non-coding elements that regulate the dynamics of gene expression, frequently, but not exclusively, via chromatin (re)modeling ([Harrow et al., 2012](#); [Maurano et al., 2012](#)). Non-coding regulatory elements include several families of non-coding RNA molecules such as long intergenic non-coding RNAs (lincRNA), and short RNA molecules, including microRNA (miRNA), short interfering RNA (siRNA), Piwi-interacting RNA (piRNA), regulatory DNA sequences, i.e. Transcription Factors, and other sequences with yet unknown functions. In addition to this network of regulatory “switches”, the genomic architecture is also characterized by low-complexity regulatory elements (e.g. LINEs, SINEs, HERVs and SVAs) that compose about 45-62 % of our genome, and consist of repeated transposable elements (TEs: DNA elements that have the ability to move, i.e., transpose, within the genome) that make humans unique in having

the largest ratio of non-coding to coding DNA, specifically in central nervous system-related genes (Taft et al., 2007; de Koning et al., 2011) (Figure 1).

Transposable Elements introduce Genetic Variability in the Human Genome.

The retrotransposition rate of still-active TEs has been estimated to range between 0.8 to 0.6 somatic L1 insertions per neuron in the human brain (Coufal et al., 2009; Evrony et al., 2012), while the transposition rate is higher in germline cells. Not surprisingly, then, cells evolved self-defense machineries to regulate the activity of TEs through epigenetic mechanisms, such as DNA methylation (TEs account for one-third of all CpG sites) and histone modification (Slotkin and Martienssen, 2007), but also by post-transcriptional mechanisms. Methylation of CpG sites results in effectively repressing the transcription of nearby genes (*cis* mechanism). Despite the machinery used by the cell to regulate TEs' activity, some TEs escape repression and generate new insertions in germline cells and during early embryonic development, as well as in somatic tissues later in life (Baillie et al., 2011; Kazazian, 2011; Lee et al., 2012), affecting the genome in different ways as explained in the next paragraphs. The human genome has accumulated about 970,000 – 1.5 Millions LINE1s (L1s), the majority of which are retrotranspositionally inactive. The still-active LINEs belong to the L1HS family (Brouha et al., 2003) and are responsible for the majority of the ongoing retrotransposition in the human populations. These variations are called *retrotransposon insertion polymorphisms* (RIPs) or *Mobile Element Insertions* (MEIs) (Ewing and Kazazian, 2010; 2011; Schmitz, 2012). At least four studies in different populations have been conducted using next-generation, i.e. short reads based, sequencing to investigate the specific content of *de novo* retrotransposition (Beck et al., 2010; Ewing and Kazazian, 2010; Huang et al., 2010; Iskow et al., 2010). The main findings were that individual genomes are highly variable in the specific locus position of individual non-reference (i.e., polymorphic) L1 insertions and that *de novo* retrotransposition were much more frequent than previously thought. However, TEs do

not transpose within the genome in an exclusive random mode: LINEs tend to re-insert into GC-poor, and SINE/Alus into GC-rich, regions ([Pavlicek et al., 2001](#)). This evidence provides a highly dynamic portrait of the genome, where individuals differ not only with respect to the presence or absence of various L1 insertions, but to the relative position of the insertion ([Lupski, 2010](#)). See Figure 1 for a classification and schematic representation of the molecular structure of TEs.

Transposable Elements induce transcriptome diversity. TEs can affect genomic integrity in many ways through *de novo* insertions and post-insertional rearrangements, resulting in deleterious mutations in the open reading frames of a gene, leading to protein disruption and/or aberrant expression ([Goodier and Kazazian, 2008](#)). Even if transposon-mediated mutagenesis is extensive in the human genome ([Cordaux et al., 2006](#)), this is not the only mechanism by which TEs can influence cellular functions. Contrasting the hypothesis that retrotransposons are disruptive and negatively affect genomic integrity ([Goodier and Kazazian, 2008](#)), is the finding in many cases that they have rather been exapted as Transcriptional Start Sites (TSSs) or enhancers ([Huda et al., 2010](#)), and significantly and positively contributed to the evolution of the human genome ([Cordaux et al., 2006](#)).

TE non-reference or *de novo* insertions may be beneficial to gene expression in inducing transcript diversity ([Cowley and Oakey, 2013](#)). TE insertions can generate alternative splice sites, causing the splicing system to include new sequences as exons or to elongate existing ones ([Schmitz and Brosius, 2011](#)). They can also introduce poly(A) signals, causing premature termination of transcription of the host gene, and also generate new transcription start sites ([Stefansson et al.](#)). All these mechanisms have a clear role in promoting transcript diversity. Furthermore, recent genome-scale studies have revealed the importance of TEs in dispersing transcription factor binding sites (TFBS). The effects of the dispersion of TE-derived TFBS are multiple. In normal tissues, they contribute to the generation of tissue-specific (e.g., brain-specific,

and brain area-specific) expression profiles ([Faulkner et al., 2009](#)), while the inappropriate activation of these TE-derived transcription factor binding sites seem to be responsible for driving ectopic gene expression, which has been implicated in human diseases ([Faulkner et al., 2009](#)). Moreover, TEs can coordinate the activation of gene expression throughout the genome, co-orchestrating the activity of functionally related genes, i.e. within the same pathway, but not necessarily physically clustered ([Britten and Davidson, 1971](#); [Lynch et al., 2015](#)). For this fundamental activity of coordinators of global patterns of gene expression, TEs have been defined as the “engineers” of transcriptional networks ([Cowley and Oakey, 2013](#)).

TEs might also impact gene expression indirectly, when the retrotransposition machinery is used by non-TE genes to retrotranspose protein-coding mRNAs, leading to the creation of a copy of the original gene. While these copies are mostly nonfunctional, sometimes they can evolve into retrogenes with novel functions. Retrogenes, when embedded in the introns of genes, have the potential to induce premature termination of transcription by causing upstream transcript polyadenylation. This particular mechanism affects only one of the two parental alleles ([Cowley and Oakey, 2013](#)), and creates a polymorphic effect similar to that shown for intronic HERVs ([Ward et al., 2013](#)). On the basis of an increasingly overwhelming body of evidence, it is reasonable to believe that the study of the “mobilome” may transform our understanding of the mechanisms of coordination responsible for gene expression throughout transcriptional networks, leading to a better understanding of the functional architecture of the normal brain, as well as, of the etiopathogenesis of many neuropsychiatric diseases ([Erwin et al., 2014](#); [Guffanti et al., 2014](#); [Insel, 2014](#)).

An impressive level of TE transcription is present both in evolutionarily conserved and accelerated genome regions. Lots of evolutionarily “old” L1s show an intense expression even when severely truncated, while also L1a other than L1HS are present in rapidly evolving regions.

For example, there are several almost intact L1PA3, L1PA2 and L1PA4s that show a high level of expression in positively swept regions. Many TEs look unique to humans compared with other primates or vertebrates, suggesting either a more recent than expected evolutionary time or representing relatively “new” polymorphic insertions in the human genome. Despite the current evidence for the relevance of HARs (Human Accelerated Regions) in shaping the human genome ([Gilad et al., 2006](#); [Pollard et al., 2006](#); [Lindblad-Toh et al., 2011](#); [Burbano et al., 2012](#)), the real modes of evolution are not yet definitively known ([Hernandez et al., 2011](#); [McLean et al., 2011](#)). It is possible that a polygenic combination of “soft” sweeps in networks of genes ([Cordaux and Batzer, 2009](#); [Erwin and Davidson, 2009](#); [Pritchard et al., 2010](#)) rather than a positive selection at single loci may be one of the major mechanisms regulating the evolution of expression ([Khaitovich et al., 2006](#); [Feschotte, 2008](#)) especially for the developmentally relevant regulatory elements ([Matlik et al., 2006](#)).

Frequently, L1s (or Alus or HERVs) that are consistently expressed overlap with transcription signals detected by the Encode consortium ([Kundaje et al., 2015](#)), including transcription factors (TF) and transcription factor binding sites (TFBS). Frequently these transcription signals also disperse within H3K4-me1 and 3 histone marks, suggesting promoter or regulatory (enhancers / insulators) functions. The pervasive pattern of TEs transcription at a whole-genome level nonetheless is not uniform: some genes are almost spared of TEs across their entire ORF, others show an extremely dense pattern of TEs across the entire gene structure, including introns, 3' and 5' UTRs.

Thus, it is becoming clear that the non-coding genome has a critical epigenetic role in controlling for the complex pattern of tissue and time specific patterns of gene expression, as well as it is also becoming more and more evident that genes do not work in isolation, but only as part of gene-networks.

We are just beginning to understand the complexity of such regulatory genome and of its epigenetic mechanisms. Epigenetics / regulatory mechanisms can work either locally – through promoters, enhancers, insulators and other element controlling for the expression of nearby genes – or distantly – through a variety of other mechanisms, like methylation-silencing gene promoters, transcription factors, short and long non-coding RNAs, transposable elements and chromatin modifiers.

Transposon-mediated epigenetic model for positive emotions

A great deal of work has been conducted on negative emotions, such as fearful behaviors, but evidence on neurobiological mechanisms underlying positive emotions is still relatively limited ([Burgdorf and Panksepp, 2006](#)). As a consequence, much of what we know on epigenetic mechanisms promoting the transduction of environmental inputs into affective states derives from the stress-fear-response model.

First, we review a working example of interaction between stress, epigenetics and emotions involving specifically Transposable Elements as “controlling elements”. It is plausible to argue that positive emotions might be mediated by the same biological mechanisms. Second, we review a working example of interaction between neuroimaging, epigenetics and emotions involving brain regions associated with emotion regulation. It is plausible to argue that positive emotions might be mediated by the same biological mechanisms reviewed in these two experimental frameworks. In the end, we discuss how transposon-mediated epigenetic control in stress-response model may provide us with an explanation for brain plasticity at the basis of positive emotions.

Transposable Elements, Stress and Emotions. Initial evidence of an epigenetic regulation of transposable elements mediating the effects of stress on neural plasticity comes from work by Hunter and colleagues ([Hunter et al., 2013](#)). In rats, acute stress induces increased hippocampal levels of the histone H3 lysine 9 trimethylation (H3K9me3), whose function is mainly associated with repression of gene expression (Hunter et al., 2009). Further research illustrated that increased levels of this epigenetic mark were specifically localized to active classes of transposable elements, namely LINEs, SINEs and ERV/LTR, leading in turn to their reduced transcriptional levels. Such a response to stress initially revealed an high hippocampal specificity, although more comprehensive investigations are needed before ruling out similar effects in other brain regions similarly known for their role in stress-sensitivity, such as amygdala, septum, bed nucleus of the stria terminalis, nucleus accumbens, and limbic thalamus. In an effort to further explain the mechanism by which stress induces epigenetic alterations in the brain, the authors examined the expression of several methyltransferases, the enzymes that allow methylation (and therefore transcriptional repression) of their molecular targets, in this case the aforementioned active classes of TEs. Up-regulation of one of these enzymes, i.e. *Suv39h2*, was found to be associated with increased binding of the glucocorticoid receptor (GR), consistent with higher expression in the hippocampus ([Hunter et al., 2012](#); [Hunter et al., 2015](#)). The GR is a known stress-related hormone-dependent transcriptional regulator previously implicated in the studies of epigenetic mediation of stress exposure, with particular emphasis on the effects of maternal care on an individual's behavior ([Meaney and Szyf, 2005](#)).

Overall, these findings supported an active role for transposable elements in the epigenetic mechanisms underlying the biological response to environmental stress exposure. The downstream effects of such epigenetic response in shaping the programming and reprogramming of neural circuitry of emotional regulation remain to be fully elucidated. In other

words, the direction in which repression of transcriptional activity of large portions of the non-coding genome is capable of influencing widespread transcriptional activity in the brain has not yet been the object of thorough investigation. Recently, work by Fasching and colleagues (Fasching et al., 2015) illustrated a potential mechanism that could explain the complex relationship between stress-mediated epigenetic regulation of transposable elements and transcriptional plasticity of neural circuitry. The authors reported that deletion of the gene *TRIM28*, a transcriptional regulator, in neural progenitor cells determines loss of H3K9me3 epigenetic mark localized on HERVs, which is ultimately leading to their transcriptional re-activation. Interestingly, re-activation of HERVs revealed exaptation properties of this class of transposable elements by influencing expression levels of nearby genes and production of long noncoding RNAs, also known for cis- and trans- regulatory functions themselves (Guttman and Rinn, 2012). Upon loss of repressive epigenetic mark, HERVs were capable of acting as promoters/enhancers inducing gene expression of other genes, potentially reconfiguring the whole transcriptional network (Feschotte, 2008). Although far from conclusive, these findings encourage further explorations of transposable elements and their exaptation into cis-regulatory elements of transcriptional activity.

Epigenetics, Brain circuits, and Emotions. Studies related to the genomic / epigenomic regulation of positive emotions are just beginning to appear in the current literature (Puglia et al., 2015). In this case, the authors provide evidence of association of higher levels of oxytocin receptor gene methylation with higher activation of brain areas involved in emotional regulation, namely amygdala, fusiform and insula and with “decreased functional coupling of amygdala with the rest of the regions involved in emotion regulation”, ultimately leading to increased anxiety.

Although preliminary, these findings support the relevance of epigenetic regulation in shaping the neural circuitry of emotional regulation.

The integration of epigenetics and brain circuitry of emotional regulation extends efforts to include epigenetics into the framework of genome-wide association studies other than simply inspecting candidate genes. Although the era of “large-scale studies of human disease-associated epigenetic variation, specifically variation in DNA methylation” has started to deliver the first wave of results (Rakyan et al., 2011), this neuro-epigenetics study offers an example of an alternative and complimentary approach to Epigenome Wide Association Studies (EWAS) while providing an additional level of functional characterization of candidate genes findings.

In parallel, refinement of our understanding of pleasure mechanisms in the brain via imaging studies ([Berridge and Kringelbach, 2015](#)) is building up a better systematic knowledge of emotional mechanisms. These investigations suggest the existence of a neural circuit common to diverse pleasures, with specific “hot-spots” in the limbic circuitry, e.g, the dorsomedial notch of the nucleus accumbens, and other subcortical regions mostly localized in the right (non-dominant) hemisphere for well-defined “hedonic” phenotypes. Two recent papers ([Berridge and Kringelbach, 2015](#); [Namburi et al., 2015](#)) are suggesting an interesting neural mechanism possibly controlling for positive emotions. Neurons from the basolateral amygdala complex (BLA) can control for either positive or negative associations (pleasure compared to fear, for example), projecting either to mesolimbic or subcortical structures, based on the specific stimuli that come from the environment. Figure 2 shows a detailed vision of the functional circuits involved in such control, with a speculative interpretation of the role that TEs can play in driving the fate of otherwise indistinguishable neurons toward responding to either positive or negative emotions.

Conclusions

We argue that the integration of the application of these neurobiological approaches suggest a major epigenetic role of regulatory mechanisms of the neural genome, hence including Transposable Elements, to explain the biological mechanisms underlying emotions. The genome responds to environmental inputs via dynamic regulation of the expression of Transposable Elements in the brain, which in turn has an impact on protein-coding mRNA transcription and/or post-translational mechanisms. These would ultimately modulate behavioral phenotypes such as positive emotions. Because of the high level of inter-individual variation of TE-derived regulatory elements in the human genome, their ability to modulate both positive and negative exposure through epigenetics mechanisms and their emerging role as regulatory elements of transcriptional activity, Transposable Elements can have a central role in co-orchestrating the dynamic neural circuitry processes underlying emotional regulation.

Figures

Figure 1. Transposable element classification and molecular structure. In this schematic representation, the genomes of different TEs are not shown to scale. LTR-transposons or human endogenous retroviruses (HERVs) include two Long Terminal Repeat sequences flanking the coding sequences of the functional polyproteins capsid (gag), protease, polymerase and reverse transcriptase (pol) and envelope (env). Non-LTR-retrotransposons include long interspersed elements (LINEs), short interspersed elements (SINEs), and SVAs (SINE-R/VNTR/ Alu-like). L1 elements consist of an intact 5' untranslated region (UTR) that functions as an internal promoter and full-length L1 mRNA of functional ORF1 (open reading frame) and ORF2 proteins, encoding for a reverse transcriptase. Alus consist of a tRNA-related region that represents the internal promoter stretch, followed by a tRNA unrelated region and a LINE-related region, which is used by the SINE element to bind LINE-encoded proteins to complete LINE-1-mediated retrotransposition. SVAs consist of four domains including a CT-rich repeat at the 5' end, commonly referred to as CT-hexamer, an Alu-like sequence, so-called for homology with two antisense Alu-like fragments, a GC-variable number tandem repeats (VNTR), whose length determines variation of the full-length SVA elements, a sequence derived from the envelope gene (env) and a SINE-R, an LTR of an extinct HERV-K10. In the DNA transposons a protein called transposase is bound by terminal inverted repeats (TIRs) that flank the TE, excises the TE out of the donor position and re-integrates it into the genome. Schematic representation has been inspired by images in (Guffanti et al., 2014).

Figure 2. Functional circuits of positive emotions. The left side panels (A,C) depict a general model of the cortico-striato-pallido-SN-VTA downstream systems in the human brain, while the right side panels (B,D) depict the specific case of the cortico-striato-pallido-SN-VTA systems that regulate positive and negative emotions. In both cases, the systems are made up of parallel,

hierarchical, interacting cortico-subcortical modules that ultimately control behavior through downstream control of skeletal, autonomic, and endocrine motor systems, but also involve recurring loops back to both neocortex and limbic cortices, primarily through ascending thalamo-cortical and monoaminergic projections (not shown). In the general forebrain cortico-striatal originating channels in **A and C**, the sensory cortices provide topographical inputs (S for sensory) to the matrix compartments of the caudate-putamen, while premotor and motor cortices innervate adjacent matrix compartments (M for motor). Non topographical inputs from limbic cortices primarily innervate the “patch” striatal compartment. Interneurons, including tonically active neurons interconnect these compartments. These compartments then project to either the internal or external segment of the globus pallidus, and or different sectors of the substantia nigra-ventral tegmental area. Further downstream convergence occurs on thalamic, hypothalamic, brainstem and spinal cord cell columns and nuclei that control movement, the autonomics, pituitary, and preganglionics. In **B and D**, a similar set of cortico-striatal systems originating in the main cortical-like nucleus of the amygdala, the baso-lateral amygdaloid nucleus (BLA), innervate striatal-like targets in the central nucleus of the amygdala (CeM) for the control of negative emotions (in red) or the medial sector of the nucleus accumbens (NAc) containing the hedonistic hot spot for the control of positive emotions (in green). These limbic striatal-like targets in turn, project to their respective ventral pallidal and SN-VTA targets, and further downstream targets that regulate motor outputs related to emotionally dominated complex adaptive behaviors. As in the case of the general forebrain schema of cortico-striatal connectivity (**A,C**), the adjacent neurons at the cortical, striatal, and pallidal subserving different functionality in the emotional circuitry (**B,D**) (e.g., positive vs. negative emotions) are connected by direct connections (e.g. between neighboring pyramidal neurons at the cortical levels, as well as and indirect, interneuron mediated connectivity. These close interneuronal interactions at each level

also allow for epigenetic tuning of the two different functional systems, in part mediated by plastic control of transposon-mediated regulation of promoters and enhancers of coding genes characteristic of each neuron at each level of the system.

Accepted Article

References

- Baillie JK, Barnett MW, Upton KR, Gerhardt DJ, Richmond TA, De Sapio F, Brennan PM, Rizzu P, Smith S, Fell M, Talbot RT, Gustincich S, Freeman TC, Mattick JS, Hume DA, Heutink P, Carninci P, Jeddeloh JA, Faulkner GJ. 2011. Somatic retrotransposition alters the genetic landscape of the human brain. *Nature* 479(7374):534-537.
- Beck CR, Collier P, Macfarlane C, Malig M, Kidd JM, Eichler EE, Badge RM, Moran JV. 2010. LINE-1 retrotransposition activity in human genomes. *Cell* 141(7):1159-1170.
- Berridge KC, Kringelbach ML. 2015. Pleasure systems in the brain. *Neuron* 86(3):646-664.
- Britten RJ, Davidson EH. 1971. Repetitive and non-repetitive DNA sequences and a speculation on the origins of evolutionary novelty. *The Quarterly review of biology* 46(2):111-138.
- Brouha B, Schustak J, Badge RM, Lutz-Prigge S, Farley AH, Moran JV, Kazazian HH, Jr. 2003. Hot L1s account for the bulk of retrotransposition in the human population. *Proceedings of the National Academy of Sciences of the United States of America* 100(9):5280-5285.
- Burbano HA, Green RE, Maricic T, Lalueza-Fox C, de la Rasilla M, Rosas A, Kelso J, Pollard KS, Lachmann M, Paabo S. 2012. Analysis of human accelerated DNA regions using archaic hominin genomes. *PLoS One* 7(3):e32877.
- Burgdorf J, Panksepp J. 2006. The neurobiology of positive emotions. *Neurosci Biobehav Rev* 30(2):173-187.
- Consortium IHGS. 2004. Finishing the euchromatic sequence of the human genome. *Nature* 431(7011):931-945.
- Coop G, Witonsky D, Di Rienzo A, Pritchard JK. 2010. Using environmental correlations to identify loci underlying local adaptation. *Genetics* 185(4):1411-1423.
- Cordaux R, Batzer MA. 2009. The impact of retrotransposons on human genome evolution. *Nature reviews Genetics* 10(10):691-703.
- Cordaux R, Hedges DJ, Herke SW, Batzer MA. 2006. Estimating the retrotransposition rate of human Alu elements. *Gene* 373:134-137.
- Coufal NG, Garcia-Perez JL, Peng GE, Yeo GW, Mu Y, Lovci MT, Morell M, O'Shea KS, Moran JV, Gage FH. 2009. L1 retrotransposition in human neural progenitor cells. *Nature* 460(7259):1127-1131.
- Cowley M, Oakey RJ. 2013. Transposable elements re-wire and fine-tune the transcriptome. *PLoS genetics* 9(1):e1003234.
- Daskalakis NP, Bagot RC, Parker KJ, Vinkers CH, de Kloet ER. 2013. The three-hit concept of vulnerability and resilience: toward understanding adaptation to early-life adversity outcome. *Psychoneuroendocrinology* 38(9):1858-1873.
- Daskalakis NP, Oitzl MS, Schachinger H, Champagne DL, de Kloet ER. 2012. Testing the cumulative stress and mismatch hypotheses of psychopathology in a rat model of early-life adversity. *Physiol Behav* 106(5):707-721.
- de Koning AP, Gu W, Castoe TA, Batzer MA, Pollock DD. 2011. Repetitive elements may comprise over two-thirds of the human genome. *PLoS genetics* 7(12):e1002384.
- Dincheva I, Drysdale AT, Hartley CA, Johnson DC, Jing D, King EC, Ra S, Gray JM, Yang R, DeGruccio AM, Huang C, Cravatt BF, Glatt CE, Hill MN, Casey BJ, Lee FS. 2015. FAAH genetic variation enhances fronto-amygdala function in mouse and human. *Nat Commun* 6:6395.
- Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, Epstein CB, Frietze S, Harrow J, Kaul R, Khatun J, Lajoie BR, Landt SG, Lee BK, Pauli F, Rosenbloom KR, Sabo P, Safi A, Sanyal A, Shores N, Simon JM, Song L, Trinklein ND, Altshuler RC, Birney E, Brown JB, Cheng C, Djebali S, Dong X, Dunham I, Ernst J, Furey TS, Gerstein M, Giardine B, Greven M,

Hardison RC, Harris RS, Herrero J, Hoffman MM, Iyer S, Kellis M, Khatun J, Kheradpour P, Kundaje A, Lassman T, Li Q, Lin X, Marinov GK, Merkel A, Mortazavi A, Parker SC, Reddy TE, Rozowsky J, Schlesinger F, Thurman RE, Wang J, Ward LD, Whitfield TW, Wilder SP, Wu W, Xi HS, Yip KY, Zhuang J, Bernstein BE, Birney E, Dunham I, Green ED, Gunter C, Snyder M, Pazin MJ, Lowdon RF, Dillon LA, Adams LB, Kelly CJ, Zhang J, Wexler JR, Green ED, Good PJ, Feingold EA, Bernstein BE, Birney E, Crawford GE, Dekker J, Elinitzki L, Farnham PJ, Gerstein M, Giddings MC, Gingeras TR, Green ED, Guigo R, Hardison RC, Hubbard TJ, Kellis M, Kent WJ, Lieb JD, Margulies EH, Myers RM, Snyder M, Starnatoyannopoulos JA, Tennebaum SA, Weng Z, White KP, Wold B, Khatun J, Yu Y, Wrobel J, Risk BA, Gunawardena HP, Kuiper HC, Maier CW, Xie L, Chen X, Giddings MC, Bernstein BE, Epstein CB, Shores N, Ernst J, Kheradpour P, Mikkelsen TS, Gillespie S, Goren A, Ram O, Zhang X, Wang L, Issner R, Coyne MJ, Durham T, Ku M, Truong T, Ward LD, Altshuler RC, Eaton ML, Kellis M, Djebali S, Davis CA, Merkel A, Dobin A, Lassmann T, Mortazavi A, Tanzer A, Lagarde J, Lin W, Schlesinger F, Xue C, Marinov GK, Khatun J, Williams BA, Zaleski C, Rozowsky J, Roder M, Kokocinski F, Abdelhamid RF, Alioto T, Antoshechkin I, Baer MT, Batut P, Bell I, Bell K, Chakraborty S, Chen X, Chrast J, Curado J, Derrien T, Drenkow J, Dumais E, Dumais J, Duttagupta R, Fastuca M, Fejes-Toth K, Ferreira P, Foissac S, Fullwood MJ, Gao H, Gonzalez D, Gordon A, Gunawardena HP, Howald C, Jha S, Johnson R, Kapranov P, King B, Kingswood C, Li G, Luo OJ, Park E, Preall JB, Presaud K, Ribeca P, Risk BA, Robyr D, Ruan X, Sammeth M, Sandu KS, Schaeffer L, See LH, Shahab A, Skancke J, Suzuki AM, Takahashi H, Tilgner H, Trout D, Walters N, Wang H, Wrobel J, Yu Y, Hayashizaki Y, Harrow J, Gerstein M, Hubbard TJ, Reymond A, Antonarakis SE, Hannon GJ, Giddings MC, Ruan Y, Wold B, Carninci P, Guigo R, Gingeras TR, Rosenbloom KR, Sloan CA, Learned K, Malladi VS, Wong MC, Barber GP, Cline MS, Dreszer TR, Heitner SG, Karolchik D, Kent WJ, Kirkup VM, Meyer LR, Long JC, Maddren M, Raney BJ, Furey TS, Song L, Grassefeder LL, Giresi PG, Lee BK, Battenhouse A, Sheffield NC, Simon JM, Showers KA, Safi A, London D, Bhinge AA, Shestak C, Schaner MR, Kim SK, Zhang ZZ, Mieczkowski PA, Mieczkowska JO, Liu Z, McDaniell RM, Ni Y, Rashid NU, Kim MJ, Adar S, Zhang Z, Wang T, Winter D, Keefe D, Birney E, Iyer VR, Lieb JD, Crawford GE, Li G, Sandhu KS, Zheng M, Wang P, Luo OJ, Shahab A, Fullwood MJ, Ruan X, Ruan Y, Myers RM, Pauli F, Williams BA, Gertz J, Marinov GK, Reddy TE, Vielmetter J, Partridge EC, Trout D, Varley KE, Gasper C, Bansal A, Pepke S, Jain P, Amrhein H, Bowling KM, Anaya M, Cross MK, King B, Muratet MA, Antoshechkin I, Newberry KM, McCue K, Nesmith AS, Fisher-Aylor KI, Pusey B, DeSalvo G, Parker SL, Balasubramanian S, Davis NS, Meadows SK, Eggleston T, Gunter C, Newberry JS, Levy SE, Absher DM, Mortazavi A, Wong WH, Wold B, Blow MJ, Visel A, Pennachio LA, Elinitzki L, Margulies EH, Parker SC, Petrykowska HM, Abyzov A, Aken B, Barrell D, Barson G, Berry A, Bignell A, Boychenko V, Bussotti G, Chrast J, Davidson C, Derrien T, Despacio-Reyes G, Diekhans M, Ezkurdia I, Frankish A, Gilbert J, Gonzalez JM, Griffiths E, Harte R, Hendrix DA, Howald C, Hunt T, Jungreis I, Kay M, Khurana E, Kokocinski F, Leng J, Lin MF, Loveland J, Lu Z, Manthravadi D, Mariotti M, Mudge J, Mukherjee G, Notredame C, Pei B, Rodriguez JM, Saunders G, Sboner A, Searle S, Sisu C, Snow C, Steward C, Tanzer A, Tapanan E, Tress ML, van Baren MJ, Walters N, Washietl S, Wilming L, Zadissa A, Zhengdong Z, Brent M, Haussler D, Kellis M, Valencia A, Gerstein M, Raymond A, Guigo R, Harrow J, Hubbard TJ, Landt SG, Fietze S, Abyzov A, Addleman N, Alexander RP, Auerbach RK, Balasubramanian S, Bettinger K, Bhardwaj N, Boyle AP, Cao AR, Cayting P, Charos A, Cheng Y, Cheng C, Eastman C, Euskirchen G, Fleming JD, Grubert F, Habegger L, Hariharan M, Harmanci A, Iyenger S, Jin VX, Karczewski KJ, Kasowski M, Lacroute P, Lam H, Larnarre-Vincent N, Leng J, Lian J, Lindahl-Allen M,

- Min R, Miotto B, Monahan H, Moqtaderi Z, Mu XJ, O'Geen H, Ouyang Z, Patacsil D, Pei B, Raha D, Ramirez L, Reed B, Rozowsky J, Sboner A, Shi M, Sisú C, Slifer T, Witt H, Wu L, Xu X, Yan KK, Yang X, Yip KY, Zhang Z, Struhl K, Weissman SM, Gerstein M, Farnham PJ, Snyder M, Tenebaum SA, Penalva LO, Doyle F, Karmakar S, Landt SG, Bhanvadia RR, Choudhury A, Domanus M, Ma L, Moran J, Patacsil D, Slifer T, Victorsen A, Yang X, Snyder M, White KP, Auer T, Centarin L, Eichenlaub M, Gruhl F, Heerman S, Hoeckendorf B, Inoue D, Kellner T, Kirchmaier S, Mueller C, Reinhardt R, Schertel L, Schneider S, Sinn R, Wittbrodt B, Wittbrodt J, Weng Z, Whitfield TW, Wang J, Collins PJ, Aldred SF, Trinklein ND, Partridge EC, Myers RM, Dekker J, Jain G, Lajoie BR, Sanyal A, Balasundaram G, Bates DL, Byron R, Canfield TK, Diegel MJ, Dunn D, Ebersol AK, Ebersol AK, Frum T, Garg K, Gist E, Hansen RS, Boatman L, Haugen E, Humbert R, Jain G, Johnson AK, Johnson EM, Kuttyavin TM, Lajoie BR, Lee K, Lotakis D, Maurano MT, Neph SJ, Neri FV, Nguyen ED, Qu H, Reynolds AP, Roach V, Rynes E, Sabo P, Sanchez ME, Sandstrom RS, Sanyal A, Shafer AO, Stergachis AB, Thomas S, Thurman RE, Vernot B, Vierstra J, Vong S, Wang H, Weaver MA, Yan Y, Zhang M, Akey JA, Bender M, Dorschner MO, Groudine M, MacCoss MJ, Navas P, Stamatoyannopoulos G, Kaul R, Dekker J, Stamatoyannopoulos JA, Dunham I, Beal K, Brazma A, Flicek P, Herrero J, Johnson N, Keefe D, Lusk M, Luscombe NM, Sobral D, Vaquerizas JM, Wilder SP, Batzoglou S, Sidow A, Hussami N, Kyriazopoulou-Panagiotopoulou S, Libbrecht MW, Schaub MA, Kundaje A, Hardison RC, Miller W, Giardine B, Harris RS, Wu W, Bickel PJ, Banfai B, Boley NP, Brown JB, Huang H, Li Q, Li JJ, Noble WS, Bilmes JA, Buske OJ, Hoffman MM, Sahu AO, Kharchenko PV, Park PJ, Baker D, Taylor J, Weng Z, Iyer S, Dong X, Greven M, Lin X, Wang J, Xi HS, Zhuang J, Gerstein M, Alexander RP, Balasubramanian S, Cheng C, Harmanici A, Lochovsky L, Min R, Mu XJ, Rozowsky J, Yan KK, Yip KY, Birney E. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* 489(7414):57-74.
- Erwin DH, Davidson EH. 2009. The evolution of hierarchical gene regulatory networks. *Nature reviews Genetics* 10(2):141-148.
- Erwin JA, Marchetto MC, Gage FH. 2014. Mobile DNA elements in the generation of diversity and complexity in the brain. *Nature reviews Neuroscience* 15(8):497-506.
- Evrory GD, Cai X, Lee E, Hills LB, Elhosary PC, Lehmann HS, Parker JJ, Atabay KD, Gilmore EC, Poduri A, Park PJ, Walsh CA. 2012. Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell* 151(3):483-496.
- Ewing AD, Kazazian HH, Jr. 2010. High-throughput sequencing reveals extensive variation in human-specific L1 content in individual human genomes. *Genome Res* 20(9):1262-1270.
- Ewing AD, Kazazian HH, Jr. 2011. Whole-genome resequencing allows detection of many rare LINE-1 insertion alleles in humans. *Genome Res* 21(6):985-990.
- Fasching L, Kapopoulou A, Sachdeva R, Petri R, Jonsson ME, Manne C, Turelli P, Jern P, Cammas F, Trono D, Jakobsson J. 2015. TRIM28 Represses Transcription of Endogenous Retroviruses in Neural Progenitor Cells. *Cell reports* 10(1):20-28.
- Faulkner GJ, Kimura Y, Daub CO, Wani S, Plessy C, Irvine KM, Schroder K, Cloonan N, Steptoe AL, Lassmann T, Waki K, Hornig N, Arakawa T, Takahashi H, Kawai J, Forrest AR, Suzuki H, Hayashizaki Y, Hume DA, Orlando V, Grimmond SM, Carninci P. 2009. The regulated retrotransposon transcriptome of mammalian cells. *Nature genetics* 41(5):563-571.
- Feschotte C. 2008. Transposable elements and the evolution of regulatory networks. *Nature reviews Genetics* 9(5):397-405.
- Gilad Y, Oshlack A, Smyth GK, Speed TP, White KP. 2006. Expression profiling in primates reveals a rapid evolution of human transcription factors. *Nature* 440(7081):242-245.

- Goodier JL, Kazazian HH, Jr. 2008. Retrotransposons revisited: the restraint and rehabilitation of parasites. *Cell* 135(1):23-35.
- Guffanti G, Gaudi S, Fallon JH, Sobell J, Potkin SG, Pato C, Macciardi F. 2014. Transposable elements and psychiatric disorders. *American journal of medical genetics Part B, Neuropsychiatric genetics : the official publication of the International Society of Psychiatric Genetics* 165B(3):201-216.
- Guttman M, Rinn JL. 2012. Modular regulatory principles of large non-coding RNAs. *Nature* 482(7385):339-346.
- Harrow J, Frankish A, Gonzalez JM, Tapanari E, Diekhans M, Kokocinski F, Aken BL, Barrell D, Zadissa A, Searle S, Barnes I, Bignell A, Boychenko V, Hunt T, Kay M, Mukherjee G, Rajan J, Despacio-Reyes G, Saunders G, Steward C, Harte R, Lin M, Howald C, Tanzer A, Derrien T, Chrast J, Walters N, Balasubramanian S, Pei B, Tress M, Rodriguez JM, Ezkurdia I, van Baren J, Brent M, Haussler D, Kellis M, Valencia A, Reymond A, Gerstein M, Guigo R, Hubbard TJ. 2012. GENCODE: the reference human genome annotation for The ENCODE Project. *Genome Res* 22(9):1760-1774.
- Hernandez RD, Kelley JL, Elyashiv E, Melton SC, Auton A, McVean G, Sella G, Przeworski M. 2011. Classic selective sweeps were rare in recent human evolution. *Science* 331(6019):920-924.
- Huang CR, Schneider AM, Lu Y, Niranjan T, Shen P, Robinson MA, Steranka JP, Valle D, Civin CI, Wang T, Wheelan SJ, Ji H, Boeke JD, Burns KH. 2010. Mobile interspersed repeats are major structural variants in the human genome. *Cell* 141(7):1171-1182.
- Huda A, Marino-Ramirez L, Jordan IK. 2010. Epigenetic histone modifications of human transposable elements: genome defense versus exaptation. *Mobile DNA* 1(1):2.
- Hunter RG, Gagnidze K, McEwen BS, Pfaff DW. 2015. Stress and the dynamic genome: Steroids, epigenetics, and the transposome. *Proceedings of the National Academy of Sciences of the United States of America* 112(22):6828-6833.
- Hunter RG, McCarthy KJ, Milne TA, Pfaff DW, McEwen BS. 2009. Regulation of hippocampal H3 histone methylation by acute and chronic stress. *Proceedings of the National Academy of Sciences of the United States of America* 106(49):20912-20917.
- Hunter RG, McEwen BS, Pfaff DW. 2013. Environmental stress and transposon transcription in the mammalian brain. *Mobile genetic elements* 3(2):e24555.
- Hunter RG, Murakami G, Dewell S, Seligsohn M, Baker ME, Datson NA, McEwen BS, Pfaff DW. 2012. Acute stress and hippocampal histone H3 lysine 9 trimethylation, a retrotransposon silencing response. *Proceedings of the National Academy of Sciences of the United States of America* 109(43):17657-17662.
- Insel TR. 2014. Brain somatic mutations: the dark matter of psychiatric genetics? *Molecular psychiatry* 19(2):156-158.
- Iskow RC, McCabe MT, Mills RE, Torene S, Pittard WS, Neuwald AF, Van Meir EG, Vertino PM, Devine SE. 2010. Natural mutagenesis of human genomes by endogenous retrotransposons. *Cell* 141(7):1253-1261.
- Kazazian HH, Jr. 2011. Mobile DNA transposition in somatic cells. *BMC biology* 9:62.
- Khaitovich P, Enard W, Lachmann M, Paabo S. 2006. Evolution of primate gene expression. *Nature reviews Genetics* 7(9):693-702.
- Kundaje A, Meuleman W, Ernst J, Bilenky M, Yen A, Heravi-Moussavi A, Kheradpour P, Zhang Z, Wang J, Ziller MJ, Amin V, Whitaker JW, Schultz MD, Ward LD, Sarkar A, Quon G, Sandstrom RS, Eaton ML, Wu YC, Pfenning AR, Wang X, Claussnitzer M, Liu Y, Coarfa C, Harris RA, Shores N, Epstein CB, Gjoneska E, Leung D, Xie W, Hawkins RD, Lister R, Hong C, Gascard P, Mungall AJ, Moore R, Chuah E, Tam A, Canfield TK, Hansen RS, Kaul R, Sabo PJ,

- Bansal MS, Carles A, Dixon JR, Farh KH, Feizi S, Karlic R, Kim AR, Kulkarni A, Li D, Lowdon R, Elliott G, Mercer TR, Neph SJ, Onuchic V, Polak P, Rajagopal N, Ray P, Sallari RC, Siebenthall KT, Sinnott-Armstrong NA, Stevens M, Thurman RE, Wu J, Zhang B, Zhou X, Beaudet AE, Boyer LA, De Jager PL, Farnham PJ, Fisher SJ, Haussler D, Jones SJ, Li W, Marra MA, McManus MT, Sunyaev S, Thomson JA, Tlsty TD, Tsai LH, Wang W, Waterland RA, Zhang MQ, Chadwick LH, Bernstein BE, Costello JF, Ecker JR, Hirst M, Meissner A, Milosavljevic A, Ren B, Stamatoyannopoulos JA, Wang T, Kellis M. 2015. Integrative analysis of 111 reference human epigenomes. *Nature* 518(7539):317-330.
- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, Funke R, Gage D, Harris K, Heaford A, Howland J, Kann L, Lehoczky J, LeVine R, McEwan P, McKernan K, Meldrim J, Mesirov JP, Miranda C, Morris W, Naylor J, Raymond C, Rosetti M, Santos R, Sheridan A, Sougnez C, Stange-Thomann N, Stojanovic N, Subramanian A, Wyman D, Rogers J, Sulston J, Ainscough R, Beck S, Bentley D, Burton J, Clee C, Carter N, Coulson A, Deadman R, Deloukas P, Dunham A, Dunham I, Durbin R, French L, Grafham D, Gregory S, Hubbard T, Humphray S, Hunt A, Jones M, Lloyd C, McMurray A, Matthews L, Mercer S, Milne S, Mullikin JC, Mungall A, Plumb R, Ross M, Shownkeen R, Sims S, Waterston RH, Wilson RK, Hillier LW, McPherson JD, Marra MA, Mardis ER, Fulton LA, Chinwalla AT, Pepin KH, Gish WR, Chissole SL, Wendl MC, Delehaunty KD, Miner TL, Delehaunty A, Kramer JB, Cook LL, Fulton RS, Johnson DL, Minx PJ, Clifton SW, Hawkins T, Branscomb E, Predki P, Richardson P, Wenning S, Slezak T, Doggett N, Cheng JF, Olsen A, Lucas S, Elkin C, Uberbacher E, Frazier M, Gibbs RA, Muzny DM, Scherer SE, Bouck JB, Sodergren EJ, Worley KC, Rives CM, Gorrell JH, Metzker ML, Naylor SL, Kucherlapati RS, Nelson DL, Weinstock GM, Sakaki Y, Fujiyama A, Hattori M, Yada T, Toyoda A, Itoh T, Kawagoe C, Watanabe H, Totoki Y, Taylor T, Weissenbach J, Heilig R, Saurin W, Artiguenave F, Brottier P, Bruls T, Pelletier E, Robert C, Wincker P, Smith DR, Doucette-Stamm L, Rubenfield M, Weinstock K, Lee HM, Dubois J, Rosenthal A, Platzer M, Nyakatura G, Taudien S, Rump A, Yang H, Yu J, Wang J, Huang G, Gu J, Hood L, Rowen L, Madan A, Qin S, Davis RW, Federspiel NA, Abola AP, Proctor MJ, Myers RM, Schmutz J, Dickson M, Grimwood J, Cox DR, Olson MV, Kaul R, Raymond C, Shimizu N, Kawasaki K, Minoshima S, Evans GA, Athanasiou M, Schultz R, Roe BA, Chen F, Pan H, Ramser J, Lehrach H, Reinhardt R, McCombie WR, de la Bastide M, Dedhia N, Blocker H, Hornischer K, Nordsiek G, Agarwala R, Aravind L, Bailey JA, Bateman A, Batzoglu S, Birney E, Bork P, Brown DG, Burge CB, Cerutti L, Chen HC, Church D, Clamp M, Copley RR, Doerks T, Eddy SR, Eichler EE, Furey TS, Galagan J, Gilbert JG, Harmon C, Hayashizaki Y, Haussler D, Hermjakob H, Hokamp K, Jang W, Johnson LS, Jones TA, Kasif S, Kasprzyk A, Kennedy S, Kent WJ, Kitts P, Koonin EV, Korf I, Kulp D, Lancet D, Lowe TM, McLysaght A, Mikkelsen T, Moran JV, Mulder N, Pollara VJ, Ponting CP, Schuler G, Schultz J, Slater G, Smit AF, Stupka E, Szustakowski J, Thierry-Mieg D, Thierry-Mieg J, Wagner L, Wallis J, Wheeler R, Williams A, Wolf YI, Wolfe KH, Yang SP, Yeh RF, Collins F, Guyer MS, Peterson J, Felsenfeld A, Wetterstrand KA, Patrinos A, Morgan MJ, de Jong P, Catanese JJ, Osoegawa K, Shizuya H, Choi S, Chen YJ. 2001. Initial sequencing and analysis of the human genome. *Nature* 409(6822):860-921.
- Lee E, Iskow R, Yang L, Gokcumen O, Haseley P, Luquette LJ, 3rd, Lohr JG, Harris CC, Ding L, Wilson RK, Wheeler DA, Gibbs RA, Kucherlapati R, Lee C, Kharchenko PV, Park PJ. 2012. Landscape of somatic retrotransposition in human cancers. *Science* 337(6097):967-971.
- Lindblad-Toh K, Garber M, Zuk O, Lin MF, Parker BJ, Washietl S, Kheradpour P, Ernst J, Jordan G, Mauceli E, Ward LD, Lowe CB, Holloway AK, Clamp M, Gnerre S, Alfoldi J, Beal K, Chang J,

- Clawson H, Cuff J, Di Palma F, Fitzgerald S, Flicek P, Guttman M, Hubisz MJ, Jaffe DB, Jungreis I, Kent WJ, Kostka D, Lara M, Martins AL, Massingham T, Moltke I, Raney BJ, Rasmussen MD, Robinson J, Stark A, Vilella AJ, Wen J, Xie X, Zody MC, Baldwin J, Bloom T, Chin CW, Heiman D, Nicol R, Nusbaum C, Young S, Wilkinson J, Worley KC, Kovar CL, Muzny DM, Gibbs RA, Cree A, Dihn HH, Fowler G, Jhangiani S, Joshi V, Lee S, Lewis LR, Nazareth LV, Okwuonu G, Santibanez J, Warren WC, Mardis ER, Weinstock GM, Wilson RK, Delehaunty K, Dooling D, Fronik C, Fulton L, Fulton B, Graves T, Minx P, Sodergren E, Birney E, Margulies EH, Herrero J, Green ED, Haussler D, Siepel A, Goldman N, Pollard KS, Pedersen JS, Lander ES, Kellis M. 2011. A high-resolution map of human evolutionary constraint using 29 mammals. *Nature* 478(7370):476-482.
- Lupski JR. 2010. Retrotransposition and structural variation in the human genome. *Cell* 141(7):1110-1112.
- Lynch VJ, Nnamani MC, Kapusta A, Brayer K, Plaza SL, Mazur EC, Emera D, Sheikh SZ, Grutzner F, Bauersachs S, Graf A, Young SL, Lieb JD, DeMayo FJ, Feschotte C, Wagner GP. 2015. Ancient Transposable Elements Transformed the Uterine Regulatory Landscape and Transcriptome during the Evolution of Mammalian Pregnancy. *Cell reports*.
- Matlik K, Redik K, Speek M. 2006. L1 antisense promoter drives tissue-specific transcription of human genes. *Journal of biomedicine & biotechnology* 2006(1):71753.
- Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, Reynolds AP, Sandstrom R, Qu H, Brody J, Shafer A, Neri F, Lee K, Kutayavin T, Stehling-Sun S, Johnson AK, Canfield TK, Giste E, Diegel M, Bates D, Hansen RS, Neph S, Sabo PJ, Heimfeld S, Raubitschek A, Ziegler S, Cotsapas C, Sotoodehnia N, Glass I, Sunyaev SR, Kaul R, Stamatoyannopoulos JA. 2012. Systematic localization of common disease-associated variation in regulatory DNA. *Science* 337(6099):1190-1195.
- McLean CY, Reno PL, Pollen AA, Bassan AI, Capellini TD, Guenther C, Indjeian VB, Lim X, Menke DB, Schaar BT, Wenger AM, Bejerano G, Kingsley DM. 2011. Human-specific loss of regulatory DNA and the evolution of human-specific traits. *Nature* 471(7337):216-219.
- Meaney MJ, Szyf M. 2005. Environmental programming of stress responses through DNA methylation: life at the interface between a dynamic environment and a fixed genome. *Dialogues Clin Neurosci* 7(2):103-123.
- Namburi P, Beyeler A, Yoroazu S, Calhoon GG, Halbert SA, Wichmann R, Holden SS, Mertens KL, Anahtar M, Felix-Ortiz AC, Wickersham IR, Gray JM, Tye KM. 2015. A circuit mechanism for differentiating positive and negative associations. *Nature* 520(7549):675-678.
- Ohno S. 1972. So much "junk" DNA in our genome. *Brookhaven Symp Biol* 23:366-370.
- Pavlicek A, Jabbari K, Paces J, Paces V, Hejnar JV, Bernardi G. 2001. Similar integration but different stability of Alus and LINEs in the human genome. *Gene* 276(1-2):39-45.
- Pollard KS, Salama SR, Lambert N, Lambot MA, Coppens S, Pedersen JS, Katzman S, King B, Onodera C, Siepel A, Kern AD, Dehay C, Igel H, Ares M, Jr., Vanderhaeghen P, Haussler D. 2006. An RNA gene expressed during cortical development evolved rapidly in humans. *Nature* 443(7108):167-172.
- Pritchard JK, Pickrell JK, Coop G. 2010. The genetics of human adaptation: hard sweeps, soft sweeps, and polygenic adaptation. *Current biology : CB* 20(4):R208-215.
- Puglia MH, Lillard TS, Morris JP, Connelly JJ. 2015. Epigenetic modification of the oxytocin receptor gene influences the perception of anger and fear in the human brain. *Proceedings of the National Academy of Sciences of the United States of America* 112(11):3308-3313.
- Rakyan VK, Down TA, Balding DJ, Beck S. 2011. Epigenome-wide association studies for common human diseases. *Nature reviews Genetics* 12(8):529-541.

- Russo SJ, Nestler EJ. 2013. The brain reward circuitry in mood disorders. *Nature reviews Neuroscience* 14(9):609-625.
- Schmidt MV. 2011. Animal models for depression and the mismatch hypothesis of disease. *Psychoneuroendocrinology* 36(3):330-338.
- Schmitz J. 2012. SINEs as driving forces in genome evolution. *Genome dynamics* 7:92-107.
- Schmitz J, Brosius J. 2011. Exonization of transposed elements: A challenge and opportunity for evolution. *Biochimie* 93(11):1928-1934.
- Schultz W. 2011. Potential vulnerabilities of neuronal reward, risk, and decision mechanisms to addictive drugs. *Neuron* 69(4):603-617.
- Slotkin RK, Martienssen R. 2007. Transposable elements and the epigenetic regulation of the genome. *Nature reviews Genetics* 8(4):272-285.
- Stefansson H, Rujescu D, Cichon S, Pietilainen OP, Ingason A, Steinberg S, Fossdal R, Sigurdsson E, Sigmundsson T, Buizer-Voskamp JE, Hansen T, Jakobsen KD, Muglia P, Francks C, Matthews PM, Gylfason A, Halldorsson BV, Gudbjartsson D, Thorgeirsson TE, Sigurdsson A, Jonasdottir A, Bjornsson A, Mattiasdottir S, Blondal T, Haraldsson M, Magnusdottir BB, Giegling I, Moller HJ, Hartmann A, Shianna KV, Ge D, Need AC, Crombie C, Fraser G, Walker N, Lonnqvist J, Suvisaari J, Tuulio-Henriksson A, Paunio T, Touloupoulou T, Bramon E, Di Forti M, Murray R, Ruggeri M, Vassos E, Tosato S, Walshe M, Li T, Vasilescu C, Muhleisen TW, Wang AG, Ullum H, Djurovic S, Melle I, Olesen J, Kiemeny LA, Franke B, Sabatti C, Freimer NB, Gulcher JR, Thorsteinsdottir U, Kong A, Andreassen OA, Ophoff RA, Georgi A, Rietschel M, Werge T, Petursson H, Goldstein DB, Nothen MM, Peltonen L, Collier DA, St Clair D, Stefansson K. 2008. Large recurrent microdeletions associated with schizophrenia. *Nature* 455(7210):232-236.
- Taft RJ, Pheasant M, Mattick JS. 2007. The relationship between non-protein-coding DNA and eukaryotic complexity. *BioEssays : news and reviews in molecular, cellular and developmental biology* 29(3):288-299.
- Ward MC, Wilson MD, Barbosa-Morais NL, Schmidt D, Stark R, Pan Q, Schwalie PC, Menon S, Lusk M, Watt S, Thybert D, Kutter C, Kirschner K, Flicek P, Blencowe BJ, Odom DT. 2013. Latent regulatory potential of human-specific repetitive elements. *Molecular cell* 49(2):262-272.
- Yoo AS, Crabtree GR. 2009. ATP-dependent chromatin remodeling in neural development. *Curr Opin Neurobiol* 19(2):120-126.
- Yoo AS, Staahl BT, Chen L, Crabtree GR. 2009. MicroRNA-mediated switching of chromatin-remodelling complexes in neural development. *Nature* 460(7255):642-646.




Figure 1. Transposable element classification and molecular structure. In this schematic representation, the genomes of different TEs are not shown to scale. LTR-transposons or human endogenous retroviruses (HERVs) include two Long Terminal Repeat sequences flanking the coding sequences of the functional polyproteins capsid (gag), protease, polymerase and reverse transcriptase (pol) and envelope (env). Non-LTR-retrotransposons include long interspersed elements (LINEs), short interspersed elements (SINEs), and SVAs (SINE-R/VNTR/ Alu-like). L1 elements consist of an intact 5' untranslated region (UTR) that functions as an internal promoter and full-length L1 mRNA of functional ORF1 (open reading frame) and ORF2 proteins, encoding for a reverse transcriptase. Alus consist of a tRNA-related region that represents the internal promoter stretch, followed by a tRNA unrelated region and a LINE-related region, which is used by the SINE element to bind LINE-encoded proteins to complete LINE-1-mediated retrotransposition. SVAs consist of four domains including a CT-rich repeat at the 5' end, commonly referred to as CT-hexamer, an Alu-like sequence, so-called for homology with two antisense Alu-like fragments, a GC-variable number tandem repeats (VNTR), whose length determines variation of the full-length SVA elements, a sequence derived from the envelope gene (env) and a SINE-R, an LTR of an extinct HERV-K10. In the DNA transposons a protein called transposase is bound by terminal inverted repeats (TIRs) that flank the TE, excises the TE out of the donor position and re-integrates it into the genome. Schematic representation has been inspired by images in (Guffanti et al., 2014).

287x203mm (300 x 300 DPI)





Figure 2. Transposon-mediated regulation of functional circuits of positive emotions. The left side panels (A,C) depict a general model of the cortico-striato-pallido-SN-VTA downstream systems in the human brain, while the right side panels (B,D) depict the specific case of the cortico-striato-pallido-SN-VTA systems that regulate positive and negative emotions. In both cases, the systems are made up of parallel, hierarchical, interacting cortico-subcortical modules that ultimately control behavior through downstream control of skeletal, autonomic, and endocrine motor systems, but also involve recurring loops back to both neocortex and limbic cortices, primarily through ascending thalamo-cortical and monoaminergic projections (not shown). In the general forebrain cortico-striatal originating channels in A and C, the sensory cortices provide topographical inputs (S for sensory) to the matrix compartments of the caudate-putamen, while premotor and motor cortices innervate adjacent matrix compartments (M for motor). Non topographical inputs from limbic cortices primarily innervate the "patch" striatal compartment. Interneurons, including tonically active neurons interconnect these compartments. These compartments then project to either the internal or external segment of the globus pallidus, and or different sectors of the substantia nigra-ventral tegmental

Accepted Article

area. Further downstream convergence occurs on thalamic, hypothalamic, brainstem and spinal cord cell columns and nuclei that control movement, the autonomics, pituitary, and preganglionics. In B and D, a similar set of cortico-striatal systems originating in the main cortical-like nucleus of the amygdala, the basolateral amygdaloid nucleus (BLA), innervate striatal-like targets in the central nucleus of the amygdala (CeM) for the control of negative emotions (in red) or the medial sector of the nucleus accumbens (NAc) containing the hedonistic hot spot for the control of positive emotions (in green). These limbic striatal-like targets in turn, project to their respective ventral pallidal and SN-VTA targets, and further downstream targets that regulate motor outputs related to emotionally dominated complex adaptive behaviors. As in the case of the general forebrain schema of cortico-striatal connectivity (A,C), the adjacent neurons at the cortical, striatal, and pallidal subserving different functionality in the emotional circuitry (B,D) (e.g., positive vs. negative emotions) are connected by direct connections (e.g. between neighboring pyramidal neurons at the cortical levels, as well as and indirect, interneuron mediated connectivity). These close interneuronal interactions at each level also allow for epigenetic tuning of the two different functional systems, in part mediated by plastic control of transposon-mediated regulation of promoters and enhancers of coding genes characteristic of each neuron at each level of the system.

171x230mm (300 x 300 DPI)



141x105mm (72 x 72 DPI)

Graphical Abstract – text

Transposable Elements (TEs) play an essential role in brain development, organizing the modular co-expression of gene regulatory networks and thus explaining, in part, the evolutionary innovations mechanisms that underlie the architecture of the human neural genome. By regulating how genes that are localized on different genomic regions can be co-expressed in a coordinated way, ultimately controlling for the proper development and function of specific brain circuits, TEs can contribute to shape our emotional and social brain.